


Kinetic, multi-ion effects, and i-e equilibration studies in ICF plasmas using multiple nuclear and xray emission histories

Particle x-ray temporal diagnostic (PXTD)




Timing precision now on the **same** diagnostic

Xray	10 ps
D3He-p	10 - 15 ps
DT-n	10 - 15 ps
DD-n	15 – 25 ps

Shot 75694


Streak of D3He and DD neutrons



2016-04-04

Shot 80705

Streak of X-rays at three different energies



H. Sio

Hong Sio, MIT

April 5th, 2016, Kinetic Workshop
Livermore, CA

Collaborators

J. Frenje, M. Gatu Johnson, F. Seguin, C. K. Li, R. D. Petrasso

Plasma Science and Fusion Center, Massachusetts Institute of Technology

J. Katz, C. Sorce, C. Stoeckl, D. Weiner, S. Regan, M. Rosenberg,

T. C. Sangster, V. Glebov, R. Betti, F. J. Marshall, W. Seka, ...

Laboratory for Laser Energetics, University of Rochester

H. Rinderknecht, P. Amendt, S. Wilks, L. Berzak-Hopkins, ...

Lawrence Livermore National Laboratory

T. Kwan, A. Le, W. Taitano, A. Simakov, L. Chacon, A. Zylstra, N. Hoffman, K. Molvig, G. Kagan
Los Alamos National Laboratory

S. Atzeni

Universita di Roma



R. Mancini

University of Nevada, Reno



Kinetic, multi-ion effects, and i-e equilibration studies in ICF plasmas using multiple nuclear and xray emission histories

- Goal of the Particle Xray Temporal Diagnostic (PXTD) is to simultaneously provide precision time-resolved data on several nuclear reactions, as well as core xray continuum (timing precision $\pm 10\text{-}20$ ps)
- Simultaneously measured nuclear and xray emission histories - $T_i(t)$ and $T_e(t)$ - enables study of i-e equilibration / Coulomb logarithm at various plasma conditions
- Three upcoming OMEGA experiments will focus on kinetic, multi-ion effects, and i-e coupling studies. Kinetic-ion / multi-ions simulation support will be provided by LANL
 - April 13th, 2016
 - August 24th, 2016
 - August 31st, 2016

Kinetic conditions in shock-driven, thin-shell implosions are similar to the shock propagation phase of hot-spot ignition experiments



Different aspect of the burn histories (burn onset, bang-time, absolute rate) are being compared to different simulations to infer kinetic and multi-ion fluid effects






$3.8 \mu\text{m SiO}_2$



When comparing single-ion-fluid to multi-ion-fluid simulations, we observe:

- ~ 40% higher DD burn rate (relative to DT rate)
- ~ 50 ps earlier DD burn onset (relative to DT burn onset)
- ~ 30 ps earlier DD bang-time (relative to DT bang-time)

Different aspect of the burn histories (burn onset, bang-time, absolute rate) are being compared to different simulations to infer kinetic and multi-ion fluid effects




When comparing single-ion-fluid to multi-ion-fluid simulations, we observe:


- ~ 40% higher DD burn rate (relative to DT rate)
- ~ 50 ps earlier DD burn onset (relative to DT burn onset)
- ~ 30 ps earlier DD bang-time (relative to DT bang-time)

Different aspect of the burn histories (burn onset, bang-time, absolute rate) are being compared to different simulations to infer kinetic and multi-ion fluid effects

Single-ion-fluid LSP – burn histories



Two-ion-fluids LSP – burn histories





$3.8 \mu\text{m SiO}_2$



When comparing single-ion-fluid to multi-ion-fluid simulations, we observe:

- ~ 40% higher DD burn rate (relative to DT rate)
- ~ 50 ps earlier DD burn onset (relative to DT burn onset)
- ~ 30 ps earlier DD bang-time (relative to DT bang-time)

Different aspect of the burn histories (burn onset, bang-time, absolute rate) are being compared to different simulations to infer kinetic and multi-ion fluid effects




$3.8 \mu\text{m SiO}_2$



When comparing single-ion-fluid to multi-ion-fluid simulations, we observe:


- ~ 40% higher DD burn rate (relative to DT rate)
- ~ 50 ps earlier DD burn onset (relative to DT burn onset)
- ~ 30 ps earlier DD bang-time (relative to DT bang-time)

In NIF exploding pusher implosions, nuclear and xray bang times are measured with two different diagnostics to a precision of ~70ps



On OMEGA, we can make x-ray and nuclear bang-time measurements with much higher precision using PXTD (10-20 ps vs. ~ 70 ps)

The new PXTD will simultaneously provide precision time-resolved data on several nuclear reactions, as well as core x-ray continuum




Timing precision on the same diagnostic

Xray	10 ps
D3He-p	10 - 15 ps
DT-n	10 - 15 ps
DD-n	15 – 25 ps

Shot 75694

Streak of D3He and DD neutrons




2016-04-04

H. Sio

Shot 80705



Streak of X-rays at three different energies



Stoeckl *et al.*, RSI 2003


Sio *et al.*, RSI 2016 (in progress)

The burn histories is recovered from the leading edge of the streak signal using a deconvolution or forward fit approach




Different aspects of the burn histories (bang-time, width, onset) can be quantitatively compared with different simulations

Single-ion-fluid simulation





Measured burn histories



When comparing burn histories measurements to single-ion-fluid , we observe:





- D^3He burn onset and bang-time are earlier relative to DT, in good agreement with simulation. Longer burn duration ($\sim 150 \text{ ps}$) than simulation ($\sim 120 \text{ ps}$)
- For this relatively hydrodynamic-like implosion (1.3 mg/cc), maybe we do expect fairly good agreement between measurements and single-fluid simulation

To understand i-e equilibration rate, we need to determine
 $\langle T_i(t) \rangle$ from the temperature-dependent ratio of two nuclear reaction histories





Making both burn history measurements on a single diagnostic eliminates cross-timing issue

To understand i-e equilibration rate, we need to determine
 $\langle T_e(t) \rangle$ from the slope of the core xray continuum




A similar approach is being used on the NIF


Shock-driven exploding pusher is a promising system to study ion-electron equilibration because shock passage primarily heats the ions



To make the best measurements the plasma temperature should be sensitive to i-e coupling, and other energy-transfer mechanisms (i.e. thermal conduction) need be well-understood




The OMEGA experiment next week will focus on measuring both nuclear and xray emission histories in kinetic and more hydrodynamic-like ICF implosions



OMEGA experiments in August will use trace tritium to measure both D³He and DT reaction histories, in addition to xray emission histories

The OMEGA experiment next week will focus on measuring both nuclear and xray emission histories in kinetic and more hydrodynamic-like ICF implosions



180um Al + 50um Ti
(xray > 6 keV)

180um Al + 600um SS
(xray > 18 keV)

1500um Al
(xray > 9 keV)

Kinetic, multi-ion effects, and i-e equilibration studies in ICF plasmas using multiple nuclear and xray emission histories

- Goal of the Particle Xray Temporal Diagnostic (PXTD) is to simultaneously provide precision time-resolved data on several nuclear reactions, as well as core xray continuum (timing precision $\pm 10\text{-}20$ ps)
- Simultaneously measured nuclear and xray emission histories - $T_i(t)$ and $T_e(t)$ - enables study of i-e equilibration / Coulomb logarithm at various plasma conditions
- Three upcoming OMEGA experiments will focus on kinetic, multi-ion effects, and i-e coupling studies. Kinetic-ion / multi-ions simulation support will be provided by LANL
 - April 13th, 2016
 - August 24th, 2016
 - August 31st, 2016